

# ON THE EXACTNESS OF ORDINARY PARTS OVER A LOCAL FIELD OF CHARACTERISTIC $p$

JULIEN HAUSEUX

ABSTRACT. Let  $G$  be a connected reductive group over a non-archimedean local field  $F$  of residue characteristic  $p$ ,  $P$  be a parabolic subgroup of  $G$ , and  $R$  be a commutative ring. When  $R$  is artinian,  $p$  is nilpotent in  $R$ , and  $\text{char}(F) = p$ , we prove that the ordinary part functor  $\text{Ord}_P$  is exact on the category of admissible smooth  $R$ -representations of  $G$ . We derive some results on Yoneda extensions between admissible smooth  $R$ -representations of  $G$ .

## 1. RESULTS

Let  $F$  be a non-archimedean local field of residue characteristic  $p$ . Let  $\mathbf{G}$  be a connected reductive algebraic  $F$ -group and  $G$  denote the topological group  $\mathbf{G}(F)$ . We let  $\mathbf{P} = \mathbf{MN}$  be a parabolic subgroup of  $\mathbf{G}$ . We write  $\bar{\mathbf{P}} = \bar{\mathbf{M}}\bar{\mathbf{N}}$  for the opposite parabolic subgroup.

Let  $R$  be a commutative ring. We write  $\text{Mod}_G^\infty(R)$  for the category of smooth  $R$ -representations of  $G$  (i.e.  $R[G]$ -modules  $\pi$  such that for all  $v \in \pi$  the stabiliser of  $v$  is open in  $G$ ) and  $R[G]$ -linear maps. It is an  $R$ -linear abelian category. When  $R$  is noetherian, we write  $\text{Mod}_G^{\text{adm}}(R)$  for the full subcategory of  $\text{Mod}_G^\infty(R)$  consisting of admissible representations (i.e. those representations  $\pi$  such that  $\pi^H$  is finitely generated over  $R$  for any open subgroup  $H$  of  $G$ ). It is closed under passing to subrepresentations and extensions, thus it is an  $R$ -linear exact subcategory, but quotients of admissible representations may not be admissible when  $\text{char}(F) = p$  (see [AHV17, Example 4.4]).

Recall the smooth parabolic induction functor  $\text{Ind}_P^G : \text{Mod}_M^\infty(R) \rightarrow \text{Mod}_G^\infty(R)$ , defined on any smooth  $R$ -representation  $\sigma$  of  $M$  as the  $R$ -module  $\text{Ind}_P^G(\sigma)$  of locally constant functions  $f : G \rightarrow \sigma$  satisfying  $f(m\bar{n}g) = m \cdot f(g)$  for all  $m \in M$ ,  $\bar{n} \in \bar{N}$ , and  $g \in G$ , endowed with the smooth action of  $G$  by right translation. It is  $R$ -linear, exact, and commutes with small direct sums. In the other direction, there is the ordinary part functor  $\text{Ord}_P : \text{Mod}_G^\infty(R) \rightarrow \text{Mod}_M^\infty(R)$  ([Eme10a, Vig16]). It is  $R$ -linear and left exact. When  $R$  is noetherian,  $\text{Ord}_P$  also commutes with small inductive limits, both functors respect admissibility, and the restriction of  $\text{Ord}_P$  to  $\text{Mod}_G^{\text{adm}}(R)$  is right adjoint to the restriction of  $\text{Ind}_P^G$  to  $\text{Mod}_M^{\text{adm}}(R)$ .

**Theorem 1.** *If  $R$  is artinian,  $p$  is nilpotent in  $R$ , and  $\text{char}(F) = p$ , then  $\text{Ord}_P$  is exact on  $\text{Mod}_G^{\text{adm}}(R)$ .*

Thus the situation is very different from the case  $\text{char}(F) = 0$  (see [Eme10b]). On the other hand if  $R$  is artinian and  $p$  is invertible in  $R$ , then  $\text{Ord}_P$  is isomorphic on  $\text{Mod}_G^{\text{adm}}(R)$  to the Jacquet functor with respect to  $P$  (i.e. the  $N$ -coinvariants) twisted by the inverse of the modulus character  $\delta_P$  of  $P$  ([AHV17, Corollary 4.19]), so that it is exact on  $\text{Mod}_G^{\text{adm}}(R)$  without any assumption on  $\text{char}(F)$ .

---

2010 *Mathematics Subject Classification.* 22E50.

*Key words and phrases.* Local fields, reductive groups, admissible smooth representations, parabolic induction, ordinary parts, extensions.

This research was partly supported by EPSRC grant EP/L025302/1.

*Remark.* Without any assumption on  $R$ ,  $\mathrm{Ind}_P^G : \mathrm{Mod}_M^\infty(R) \rightarrow \mathrm{Mod}_G^\infty(R)$  admits a left adjoint  $L_P^G : \mathrm{Mod}_G^\infty(R) \rightarrow \mathrm{Mod}_M^\infty(R)$  (the Jacquet functor with respect to  $P$ ) and a right adjoint  $R_P^G : \mathrm{Mod}_G^\infty(R) \rightarrow \mathrm{Mod}_M^\infty(R)$  ([Vig16, Proposition 4.2]). If  $R$  is noetherian and  $p$  is nilpotent in  $R$ , then  $R_P^G$  is isomorphic to  $\mathrm{Ord}_P$  on  $\mathrm{Mod}_G^{\mathrm{adm}}(R)$  ([AHV17, Corollary 4.13]). Thus under the assumptions of Theorem 1,  $R_P^G$  is exact on  $\mathrm{Mod}_G^{\mathrm{adm}}(R)$ . On the other hand if  $R$  is noetherian and  $p$  is invertible in  $R$ , then  $R_P^G$  is expected to be isomorphic to  $\delta_P L_P^G$  ('second adjointness'), and this is proved in the following cases: when  $R$  is the field of complex numbers ([Ber87]) or an algebraically closed field of characteristic  $\ell \neq p$  ([Vig96, II.3.8 2]); when  $\mathbf{G}$  is a Levi subgroup of a general linear group or a classical group with  $p \neq 2$  ([Dat09, Théorème 1.5]); when  $\mathbf{P}$  is a minimal parabolic subgroup of  $\mathbf{G}$  (see also [Dat09]). In particular,  $L_P^G$  and  $R_P^G$  are exact in all these cases.

*Question.* Are  $L_P^G$  and  $R_P^G$  exact when  $R$  is noetherian,  $p$  is nilpotent in  $R$ , and  $\mathrm{char}(F) = p$ ?

We derive from Theorem 1 some results on Yoneda extensions between admissible  $R$ -representations of  $G$ . We compute the  $R$ -modules  $\mathrm{Ext}_G^\bullet$  in  $\mathrm{Mod}_G^{\mathrm{adm}}(R)$ .

**Corollary 2.** *Assume  $R$  artinian,  $p$  nilpotent in  $R$ , and  $\mathrm{char}(F) = p$ . Let  $\sigma$  and  $\pi$  be admissible  $R$ -representations of  $M$  and  $G$  respectively. For all  $n \geq 0$ , there is a natural  $R$ -linear isomorphism*

$$\mathrm{Ext}_M^n(\sigma, \mathrm{Ord}_P(\pi)) \xrightarrow{\sim} \mathrm{Ext}_G^n(\mathrm{Ind}_P^G(\sigma), \pi).$$

This is in contrast with the case  $\mathrm{char}(F) = 0$  (see [Hau16b]). A direct consequence of Corollary 2 is that under the same assumptions,  $\mathrm{Ind}_P^G$  induces an isomorphism between the  $\mathrm{Ext}^n$  for all  $n \geq 0$  (Corollary 5). When  $R = C$  is an algebraically closed field of characteristic  $p$  and  $\mathrm{char}(F) = p$ , we determine the extensions between certain irreducible admissible  $C$ -representations of  $G$  using the classification of [AHHV17] (Proposition 6). In particular, we prove that there exists no non-split extension of an irreducible admissible  $C$ -representation  $\pi$  of  $G$  by a supersingular  $C$ -representation of  $G$  when  $\pi$  is not the extension to  $G$  of a supersingular representation of a Levi subgroup of  $G$  (Corollary 7). When  $\mathbf{G} = \mathrm{GL}_2$ , this was first proved by Hu ([Hu17, Theorem A.2]).

**Acknowledgements.** The author would like to thank F. Herzig and M.-F. Vignéras for several comments on the first version of this paper.

## 2. PROOFS

**2.1. Hecke action.** In this subsection,  $\mathbf{M}$  denotes a linear algebraic  $F$ -group and  $\mathbf{N}$  denotes a split unipotent algebraic  $F$ -group (see [CGP15, Appendix B]) endowed with an action of  $\mathbf{M}$  that we identify with the conjugation in  $\mathbf{M} \ltimes \mathbf{N}$ . We fix an open submonoid  $M^+$  of  $M$  and a compact open subgroup  $N_0$  of  $N$  stable under conjugation by  $M^+$ .

If  $\pi$  is a smooth  $R$ -representation of  $M^+ \ltimes N_0$ , then the  $R$ -modules  $\mathbf{H}^\bullet(N_0, \pi)$ , computed using the homogeneous cochain complex  $\mathbf{C}^\bullet(N_0, \pi)$  (see [NSW08, § I.2]), are naturally endowed with the Hecke action of  $M^+$ , defined as the composite

$$\mathbf{H}^\bullet(N_0, \pi) \xrightarrow{m} \mathbf{H}^\bullet(mN_0m^{-1}, \pi) \xrightarrow{\mathrm{cor}} \mathbf{H}^\bullet(N_0, \pi)$$

for all  $m \in M^+$ . At the level of cochains, this action is explicitly given as follows (see [NSW08, § I.5]). We fix a set of representatives  $\overline{N_0/mN_0m^{-1}} \subseteq N_0$  of the left cosets  $N_0/mN_0m^{-1}$  and we write  $n \mapsto \tilde{n}$  for the projection  $N_0 \rightarrow \overline{N_0/mN_0m^{-1}}$ .

For  $\phi \in C^k(N_0, \pi)$ , we have

$$(1) \quad (m \cdot \phi)(n_0, \dots, n_k) = \sum_{\bar{n} \in \overline{N_0/mN_0m^{-1}}} \bar{n}m \cdot \phi(m^{-1}\bar{n}^{-1}n_0n_0^{-1}\bar{n}m, \dots, m^{-1}\bar{n}^{-1}n_kn_k^{-1}\bar{n}m)$$

for all  $(n_0, \dots, n_k) \in N_0^{k+1}$ .

**Lemma 3.** *Assume  $p$  nilpotent in  $R$  and  $\text{char}(F) = p$ . Let  $\pi$  be a smooth  $R$ -representation of  $M^+ \times N_0$  and  $m \in M^+$ . If the Hecke action  $h_{N_0, m}$  of  $m$  on  $\pi^{N_0}$  is locally nilpotent (i.e. for all  $v \in \pi^{N_0}$  there exists  $r \geq 0$  such that  $h_{N_0, m}^r(v) = 0$ ), then the Hecke action of  $m$  on  $H^k(N_0, \pi)$  is locally nilpotent for all  $k \geq 0$ .*

*Proof.* First, we prove the lemma when  $pR = 0$ , i.e.  $R$  is a commutative  $\mathbb{F}_p$ -algebra. We assume that the Hecke action of  $m$  on  $\pi^{N_0}$  is locally nilpotent and we prove the result together with the following fact: there exists a set of representatives  $\overline{N_0/mN_0m^{-1}} \subseteq N_0$  of the left cosets  $N_0/mN_0m^{-1}$  such that the action of

$$S := \sum_{\bar{n} \in \overline{N_0/mN_0m^{-1}}} \bar{n}m \in \mathbb{F}_p[M^+ \times N_0]$$

on  $\pi$  is locally nilpotent.

We proceed by induction on the dimension of  $\mathbf{N}$  (recall that  $\mathbf{N}$  is split so that it is smooth and connected). If  $\mathbf{N} = 1$ , then the (Hecke) action of  $m$  on  $\pi^{N_0} = \pi$  is locally nilpotent by assumption, so that the result and the fact are trivially true. Assume  $\mathbf{N} \neq 1$  and that the result and the fact are true for groups of smaller dimension. Since  $\mathbf{N}$  is split, it admits a non-trivial central subgroup isomorphic to the additive group. We let  $\mathbf{N}'$  be the subgroup of  $\mathbf{N}$  generated by all such subgroups. It is a non-trivial vector group (i.e. isomorphic to a direct product of copies of the additive group) which is central (hence normal) in  $\mathbf{N}$  and stable under conjugation by  $\mathbf{M}$  (since it is a characteristic subgroup of  $\mathbf{N}$ ). We set  $\mathbf{N}'' := \mathbf{N}/\mathbf{N}'$ . It is a split unipotent algebraic  $F$ -group endowed with the induced action of  $\mathbf{M}$  and  $\dim(\mathbf{N}'') < \dim(\mathbf{N})$ . Since  $\mathbf{N}'$  is split, we have  $N'' = N'/N'$ . We write  $N'_0$  and  $N''_0$  for the compact open subgroups  $N' \cap N_0$  and  $N_0/N'_0$  of  $N'$  and  $N''$  respectively. They are stable under conjugation by  $M^+$ . We fix a set-theoretic section  $[-] : N''_0 \hookrightarrow N_0$ .

Since  $\mathbf{N}'$  is commutative and  $p$ -torsion,  $N'_0$  is a compact  $\mathbb{F}_p$ -vector space. Thus for any open subgroup  $N'_1$  of  $N'_0$ , the short exact sequence of compact  $\mathbb{F}_p$ -vector spaces

$$0 \rightarrow N'_1 \rightarrow N'_0 \rightarrow N'_0/N'_1 \rightarrow 0$$

splits. Indeed, it admits an  $\mathbb{F}_p$ -linear splitting (since  $\mathbb{F}_p$  is a field) which is automatically continuous (since  $N'_0/N'_1$  is discrete). In particular with  $N'_1 = mN'_0m^{-1}$ , we may and do fix a section  $N'_0/mN'_0m^{-1} \hookrightarrow N'_0$ . We write  $\overline{N'_0/mN'_0m^{-1}}$  for its image, so that  $N'_0 = \overline{N'_0/mN'_0m^{-1}} \times mN'_0m^{-1}$ , and  $n' \mapsto \bar{n}'$  for the projection  $N'_0 \rightarrow \overline{N'_0/mN'_0m^{-1}}$ . We set

$$S' := \sum_{\bar{n}' \in \overline{N'_0/mN'_0m^{-1}}} \bar{n}'m \in \mathbb{F}_p[M^+ \times N'_0].$$

For all  $n'_0 \in N'_0$ , we have  $n'_0 = \bar{n}'_0(\bar{n}'_0^{-1}n'_0)$  with  $\bar{n}'_0^{-1}n'_0 \in mN'_0m^{-1}$ , thus

$$n'_0S' = \sum_{\bar{n}' \in \overline{N'_0/mN'_0m^{-1}}} (\bar{n}'_0\bar{n}')m(m^{-1}(\bar{n}'_0^{-1}n'_0)m) = S'(m^{-1}(\bar{n}'_0^{-1}n'_0)m)$$

with  $m^{-1}(\bar{n}'_0^{-1}n'_0)m \in N'_0$  (in the first equality we use the fact that  $N'_0$  is commutative and in the second one we use the fact that  $\overline{N'_0/mN'_0m^{-1}}$  is a group). Therefore, there is an inclusion  $\mathbb{F}_p[N'_0]S' \subseteq S'\mathbb{F}_p[N'_0]$ .

The  $R$ -module  $\pi^{N'_0}$ , endowed with the induced action of  $N''_0$  and the Hecke action of  $M^+$  with respect to  $N'_0$ , is a smooth  $R$ -representation of  $M^+ \times N''_0$  (see the proof of [Hau16a, Lemme 3.2.1] in degree 0). On  $\pi^{N'_0}$ , the Hecke action of  $m$  with respect to  $N'_0$  coincides with the action of  $S'$  by definition. On  $(\pi^{N'_0})^{N''_0} = \pi^{N_0}$ , the Hecke action of  $m$  with respect to  $N''_0$  coincides with the Hecke action of  $m$  with respect to  $N_0$  (see the proof of [Hau16a, Lemme 3.2.2]) which is locally nilpotent by assumption. Thus by the induction hypothesis, there exists a set of representatives  $\overline{N''_0/mN''_0m^{-1}} \subseteq N''_0$  of the left cosets  $N''_0/mN''_0m^{-1}$  such that the action of

$$S := \sum_{\bar{n}'' \in \overline{N''_0/mN''_0m^{-1}}} [\bar{n}'']S' \in \mathbb{F}_p[M^+ \times N_0]$$

on  $\pi^{N'_0}$  is locally nilpotent. Moreover, there is an inclusion  $\mathbb{F}_p[N'_0]S \subseteq S\mathbb{F}_p[N'_0]$  (because  $N'_0$  is central in  $N_0$  and  $\mathbb{F}_p[N'_0]S' \subseteq S'\mathbb{F}_p[N'_0]$ ).

We prove the fact. By [Hau16c, Lemme 2.1],

$$\overline{N_0/mN_0m^{-1}} := \{[\bar{n}'']\bar{n}' : \bar{n}'' \in \overline{N''_0/mN''_0m^{-1}}, \bar{n}' \in \overline{N'_0/mN'_0m^{-1}}\} \subseteq N_0$$

is a set of representatives of the left cosets  $N_0/mN_0m^{-1}$ , and by definition

$$S = \sum_{\bar{n} \in \overline{N_0/mN_0m^{-1}}} \bar{n}m.$$

We prove that the action of  $S$  on  $\pi$  is locally nilpotent. We proceed as in the proof of [Hu12, Théorème 5.1 (i)]. Let  $v \in \pi$  and set  $\pi_r := \mathbb{F}_p[N'_0] \cdot (S^r \cdot v)$  for all  $r \geq 0$ . Since  $\mathbb{F}_p[N'_0]S \subseteq S\mathbb{F}_p[N'_0]$ , we have  $\pi_{r+1} \subseteq S \cdot \pi_r$  for all  $r \geq 0$ . Since  $N'_0$  is compact, we have  $\dim_{\mathbb{F}_p}(\pi_r) < \infty$  for all  $r \geq 0$ . If  $S^r \cdot v \neq 0$ , i.e.  $\pi_r \neq 0$ , for some  $r \geq 0$ , then  $\pi_r^{N'_0} \neq 0$  (because  $N'_0$  is a pro- $p$  group and  $\pi_r$  is a non-zero  $\mathbb{F}_p$ -vector space) so that  $\dim_{\mathbb{F}_p}(S \cdot \pi_r) < \dim_{\mathbb{F}_p} \pi_r$  (because the action of  $S$  on  $\pi^{N'_0}$  is locally nilpotent). Therefore  $\pi_r = 0$ , i.e.  $S^r \cdot v = 0$ , for all  $r \geq \dim_{\mathbb{F}_p}(\pi_0)$ .

We prove the result. The  $R$ -modules  $\mathbf{H}^\bullet(N'_0, \pi)$ , endowed with the induced action of  $N''_0$  and the Hecke action of  $M^+$ , are smooth  $R$ -representations of  $M^+ \times N''_0$  (see the proof of [Hau16a, Lemme 3.2.1]<sup>1</sup>). At the level of cochains, the actions of  $n'' \in N''_0$  and  $m$  are explicitly given as follows. For  $\phi \in C^j(N'_0, \pi)$ , we have

$$(2) \quad (n'' \cdot \phi)(n'_0, \dots, n'_j) = [n''] \cdot \phi(n'_0, \dots, n'_j)$$

$$(3) \quad (m \cdot \phi)(n'_0, \dots, n'_j) = S' \cdot \phi(m^{-1}n'_0\bar{n}'_0^{-1}m, \dots, m^{-1}n'_j\bar{n}'_j^{-1}m)$$

for all  $(n'_0, \dots, n'_j) \in N'^{j+1}_0$  (for (2) we use the fact that  $N'_0$  is central in  $N_0$ , for (3) we use (1) and the fact that  $n' \mapsto \bar{n}'$  is a group homomorphism  $N'_0 \rightarrow \overline{N'_0/mN'_0m^{-1}}$ ). Using (2) and (3), we can give explicitly the Hecke action of  $m$  on  $\mathbf{H}^\bullet(N'_0, \pi)^{N''_0}$  at the level of cochains as follows. For  $\phi \in C^j(N'_0, \pi)$ , we have

$$(m \cdot \phi)(n'_0, \dots, n'_j) = S \cdot \phi(m^{-1}n'_0\bar{n}'_0^{-1}m, \dots, m^{-1}n'_j\bar{n}'_j^{-1}m)$$

for all  $(n'_0, \dots, n'_j) \in N'^{j+1}_0$ . Since the action of  $S$  on  $\pi$  is locally nilpotent and the image of a locally constant cochain is finite by compactness of  $N'_0$ , we deduce that the Hecke action of  $m$  on  $\mathbf{H}^j(N'_0, \pi)^{N''_0}$  is locally nilpotent for all  $j \geq 0$ . Thus the Hecke action of  $m$  on  $\mathbf{H}^i(N''_0, \mathbf{H}^j(N'_0, \pi))$  is locally nilpotent for all  $i, j \geq 0$  by the induction hypothesis. We conclude using the spectral sequence of smooth  $R$ -representations of  $M^+$

$$\mathbf{H}^i(N''_0, \mathbf{H}^j(N'_0, \pi)) \Rightarrow \mathbf{H}^{i+j}(N_0, \pi)$$

(see the proof of [Hau16a, Proposition 3.2.3] and footnote 1).

<sup>1</sup>We do not know whether [Eme10b, Proposition 2.1.11] holds true when  $\text{char}(F) = p$ , but [Hau16a, Lemme 3.1.1] does and any injective object of  $\text{Mod}_{M^+ \times N_0}^\infty(R)$  is still  $N_0$ -acyclic.

Now, we prove the lemma without assuming  $pR = 0$ . We proceed by induction on the degree of nilpotency  $r$  of  $p$  in  $R$ . If  $r \leq 1$ , then the lemma is already proved. We assume  $r > 1$  and that we know the lemma for rings in which the degree of nilpotency of  $p$  is  $r - 1$ . There is a short exact sequence of smooth  $R$ -representations of  $M^+ \ltimes N_0$

$$0 \rightarrow p\pi \rightarrow \pi \rightarrow \pi/p\pi \rightarrow 0.$$

Taking the  $N_0$ -cohomology yields a long exact sequence of smooth  $R$ -representations of  $M^+$

$$(4) \quad 0 \rightarrow (p\pi)^{N_0} \rightarrow \pi^{N_0} \rightarrow (\pi/p\pi)^{N_0} \rightarrow H^1(N_0, p\pi) \rightarrow \cdots.$$

If the Hecke action of  $m$  on  $\pi^{N_0}$  is locally nilpotent, then the Hecke action of  $m$  on  $(p\pi)^{N_0}$  is also locally nilpotent so that the Hecke action of  $m$  on  $H^k(N_0, p\pi)$  is locally nilpotent for all  $k \geq 0$  by the induction hypothesis (since  $p\pi$  is an  $R/p^{r-1}R$ -module). Using (4), we deduce that the Hecke action of  $m$  on  $(\pi/p\pi)^{N_0}$  is also locally nilpotent so that the Hecke action of  $m$  on  $H^k(N_0, \pi/p\pi)$  is locally nilpotent for all  $k \geq 0$  (since  $\pi/p\pi$  is an  $\mathbb{F}_p$ -vector space). Using again (4), we conclude that the Hecke action of  $m$  on  $H^k(N_0, \pi)$  is locally nilpotent for all  $k \geq 0$ .  $\square$

**2.2. Proof of the main result.** We fix a compact open subgroup  $N_0$  of  $N$  and we let  $M^+$  be the open submonoid of  $M$  consisting of those elements  $m$  contracting  $N_0$  (i.e.  $mN_0m^{-1} \subseteq N_0$ ). We let  $\mathbf{Z}_M$  denote the centre of  $\mathbf{M}$  and we set  $Z_M^+ := Z_M \cap M^+$ . We fix an element  $z \in Z_M^+$  strictly contracting  $N_0$  (i.e.  $\cap_{r \geq 0} z^r N_0 z^{-r} = 1$ ).

Recall that the ordinary part of a smooth  $R$ -representation  $\pi$  of  $P$  is the smooth  $R$ -representation of  $M$

$$\text{Ord}_P(\pi) := (\text{Ind}_{M^+}^M(\pi^{N_0}))^{Z_M^{-1}\text{-fin}}$$

where  $\text{Ind}_{M^+}^M(\pi^{N_0})$  is defined as the  $R$ -module of functions  $f : M \rightarrow \pi^{N_0}$  such that  $f(mm') = m \cdot f(m')$  for all  $m \in M^+$  and  $m' \in M$ , endowed with the action of  $M$  by right translation, and the superscript  $Z_M^{-1}\text{-fin}$  denotes the subrepresentation consisting of locally  $Z_M$ -finite elements (i.e. those elements  $f$  such that  $R[Z_M] \cdot f$  is contained in a finitely generated  $R$ -submodule). The action of  $M$  on the latter is smooth by [Vig16, Remark 7.6]. If  $R$  is artinian and  $\pi^{N_0}$  is locally  $Z_M^+$ -finite (i.e. it may be written as the union of finitely generated  $Z_M^+$ -invariant  $R$ -submodules), then there is a natural  $R$ -linear isomorphism

$$(5) \quad \text{Ord}_P(\pi) \xrightarrow{\sim} R[z^{\pm 1}] \otimes_{R[z]} \pi^{N_0}$$

(cf. [Eme10b, Lemma 3.2.1 (1)], whose proof also works when  $\text{char}(F) = p$  and over any artinian ring).

If  $\sigma$  is a smooth  $R$ -representation of  $M$ , then the  $R$ -module  $\mathcal{C}_c^\infty(N, \sigma)$  of locally constant functions  $f : N \rightarrow \sigma$  with compact support, endowed with the action of  $N$  by right translation and the action of  $M$  given by  $(m \cdot f) : n \mapsto m \cdot f(m^{-1}nm)$  for all  $m \in M$ , is a smooth  $R$ -representation of  $P$ . Thus we obtain a functor  $\mathcal{C}_c^\infty(N, -) : \text{Mod}_M^\infty(R) \rightarrow \text{Mod}_P^\infty(R)$ . It is  $R$ -linear, exact, and commutes with small direct sums. The results of [Eme10a, § 4.2] hold true when  $\text{char}(F) = p$  and over any ring, thus the functors

$$\begin{aligned} \mathcal{C}_c^\infty(N, -) : \text{Mod}_M^\infty(R)^{Z_M^{-1}\text{-fin}} &\rightarrow \text{Mod}_P^\infty(R) \\ \text{Ord}_P : \text{Mod}_P^\infty(R) &\rightarrow \text{Mod}_M^\infty(R)^{Z_M^{-1}\text{-fin}} \end{aligned}$$

are adjoint and the unit of the adjunction is an isomorphism.

**Lemma 4.** *Assume  $R$  artinian,  $p$  nilpotent in  $R$ , and  $\text{char}(F) = p$ . Let  $\pi$  be a smooth  $R$ -representation of  $P$ . If  $\pi^{N_0}$  is locally  $Z_M^+$ -finite, then the Hecke action of  $z$  on  $H^k(N_0, \pi)$  is locally nilpotent for all  $k \geq 1$ .*

*Proof.* We set  $\sigma := \text{Ord}_P(\pi)$ . The counit of the adjunction between  $\mathcal{C}_c^\infty(N, -)$  and  $\text{Ord}_P$  induces a natural morphism of smooth  $R$ -representations of  $P$

$$(6) \quad \mathcal{C}_c^\infty(N, \sigma) \rightarrow \pi.$$

Taking the  $N_0$ -invariants yields a morphism of smooth  $R$ -representations of  $M^+$

$$(7) \quad \mathcal{C}_c^\infty(N, \sigma)^{N_0} \rightarrow \pi^{N_0}.$$

By definition,  $\sigma$  is locally  $Z_M$ -finite so it may be written as the union of finitely generated  $Z_M$ -invariant  $R$ -submodules  $(\sigma_i)_{i \in I}$ . Thus  $\mathcal{C}_c^\infty(N, \sigma)^{N_0}$  is the union of the finitely generated  $Z_M^+$ -invariant  $R$ -submodules  $(\mathcal{C}^\infty(z^{-r}N_0z^r, \sigma_i)^{N_0})_{r \geq 0, i \in I}$ , so it is locally  $Z_M^+$ -finite. By assumption,  $\pi^{N_0}$  is also locally  $Z_M^+$ -finite. Therefore, using (5) and its analogue with  $\mathcal{C}_c^\infty(N, \sigma)$  instead of  $\pi$ , the localisation with respect to  $z$  of (7) is the natural morphism of smooth  $R$ -representations of  $M$

$$\text{Ord}_P(\mathcal{C}_c^\infty(N, \sigma)) \rightarrow \text{Ord}_P(\pi)$$

induced by applying the functor  $\text{Ord}_P$  to (6), and it is an isomorphism since the unit of the adjunction between  $\mathcal{C}_c^\infty(N, -)$  and  $\text{Ord}_P$  is an isomorphism.

Let  $\kappa$  (resp.  $\iota$ ) be the kernel (resp. image) of (6), hence two short exact sequences of smooth  $R$ -representations of  $P$

$$(8) \quad 0 \rightarrow \kappa \rightarrow \mathcal{C}_c^\infty(N, \sigma) \rightarrow \iota \rightarrow 0$$

$$(9) \quad 0 \rightarrow \iota \rightarrow \pi \rightarrow \pi/\iota \rightarrow 0$$

such that the third arrow of (8) and the second arrow of (9) fit into a commutative diagram of smooth  $R$ -representations of  $P$

$$\begin{array}{ccc} \mathcal{C}_c^\infty(N, \sigma) & \xrightarrow{\quad} & \pi \\ & \searrow & \nearrow \\ & \iota & \end{array}$$

whose upper arrow is (6). Taking the  $N_0$ -invariants yields a commutative diagram of smooth  $R$ -representations of  $M^+$

$$\begin{array}{ccc} \mathcal{C}_c^\infty(N, \sigma)^{N_0} & \xrightarrow{\quad} & \pi^{N_0} \\ & \searrow & \nearrow \\ & \iota^{N_0} & \end{array}$$

whose upper arrow is (7). Since the localisation with respect to  $z$  of the latter is an isomorphism, the localisation with respect to  $z$  of the injection  $\iota^{N_0} \hookrightarrow \pi^{N_0}$  is surjective, thus it is an isomorphism (as it is also injective by exactness of localisation). Therefore the localisation with respect to  $z$  of the morphism  $\mathcal{C}_c^\infty(N, \sigma)^{N_0} \rightarrow \iota^{N_0}$  is an isomorphism.

Since  $\mathcal{C}_c^\infty(N, \sigma) \cong \bigoplus_{n \in N/N_0} \mathcal{C}^\infty(nN_0, \sigma)$  as a smooth  $R$ -representation of  $N_0$ , it is  $N_0$ -acyclic (see [NSW08, § I.3]). Thus the long exact sequence of  $N_0$ -cohomology induced by (8) yields an exact sequence of smooth  $R$ -representations of  $M^+$

$$(10) \quad 0 \rightarrow \kappa^{N_0} \rightarrow \mathcal{C}_c^\infty(N, \sigma)^{N_0} \rightarrow \iota^{N_0} \rightarrow \mathbf{H}^1(N_0, \kappa) \rightarrow 0$$

and an isomorphism of smooth  $R$ -representations of  $M^+$

$$(11) \quad \mathbf{H}^k(N_0, \iota) \xrightarrow{\sim} \mathbf{H}^{k+1}(N_0, \kappa)$$

for all  $k \geq 1$ . Since the localisation with respect to  $z$  of the third arrow of (10) is an isomorphism, the Hecke action of  $z$  on  $\kappa^{N_0}$  is locally nilpotent. Thus the Hecke action of  $z$  on  $\mathbf{H}^k(N_0, \kappa)$  is locally nilpotent for all  $k \geq 0$  by Lemma 3. Using (11), we deduce that the Hecke action of  $z$  on  $\mathbf{H}^k(N_0, \iota)$  is locally nilpotent for all  $k \geq 1$ .

Taking the  $N_0$ -cohomology of (9) yields a long exact sequence of smooth  $R$ -representations of  $M^+$

$$(12) \quad 0 \rightarrow \iota^{N_0} \rightarrow \pi^{N_0} \rightarrow (\pi/\iota)^{N_0} \rightarrow H^1(N_0, \iota) \rightarrow \cdots$$

Since the localisation with respect to  $z$  of the second arrow is an isomorphism and the Hecke action of  $z$  on  $H^1(N_0, \iota)$  is locally nilpotent, the Hecke action of  $z$  on  $(\pi/\iota)^{N_0}$  is locally nilpotent. Thus the Hecke action of  $z$  on  $H^k(N_0, \pi/\iota)$  is locally nilpotent for all  $k \geq 0$  by Lemma 3. We conclude using (12) and the fact that the Hecke action of  $z$  on  $H^k(N_0, \iota)$  is locally nilpotent for all  $k \geq 1$ .  $\square$

*Proof of Theorem 1.* Assume  $R$  artinian,  $p$  nilpotent in  $R$ , and  $\text{char}(F) = p$ . Let

$$(13) \quad 0 \rightarrow \pi_1 \rightarrow \pi_2 \rightarrow \pi_3 \rightarrow 0$$

be a short exact sequence of admissible  $R$ -representations of  $G$ . Taking the  $N_0$ -invariants yields an exact sequence of smooth  $R$ -representations of  $M^+$

$$(14) \quad 0 \rightarrow \pi_1^{N_0} \rightarrow \pi_2^{N_0} \rightarrow \pi_3^{N_0} \rightarrow H^1(N_0, \pi_1).$$

The terms  $\pi_1^{N_0}, \pi_2^{N_0}, \pi_3^{N_0}$  are locally  $Z_M^+$ -finite (cf. [Eme10b, Theorem 3.4.7 (1)], whose proof in degree 0 also works when  $\text{char}(F) = p$  and over any noetherian ring) and the Hecke action of  $z$  on  $H^1(N_0, \pi_1)$  is locally nilpotent by Lemma 4. Therefore, using (5), the localisation with respect to  $z$  of (14) is the short sequence of admissible  $R$ -representations of  $M$

$$0 \rightarrow \text{Ord}_P(\pi_1) \rightarrow \text{Ord}_P(\pi_2) \rightarrow \text{Ord}_P(\pi_3) \rightarrow 0$$

induced by applying the functor  $\text{Ord}_P$  to (13), and it is exact by exactness of localisation.  $\square$

**2.3. Results on extensions.** We assume  $R$  noetherian. The  $R$ -linear category  $\text{Mod}_G^{\text{adm}}(R)$  is not abelian in general, but merely exact in the sense of Quillen ([Qui73]). An exact sequence of admissible  $R$ -representations of  $G$  is an exact sequence of smooth  $R$ -representations of  $G$

$$\cdots \rightarrow \pi_{n-1} \rightarrow \pi_n \rightarrow \pi_{n+1} \rightarrow \cdots$$

such that the kernel and the cokernel of every arrow are admissible. In particular, each term of the sequence is also admissible.

For  $n \geq 0$  and  $\pi, \pi'$  two admissible  $R$ -representations of  $G$ , we let  $\text{Ext}_G^n(\pi', \pi)$  denote the  $R$ -module of  $n$ -fold Yoneda extensions ([Yon60]) of  $\pi'$  by  $\pi$  in  $\text{Mod}_G^{\text{adm}}(R)$ , defined as equivalence classes of exact sequences

$$0 \rightarrow \pi \rightarrow \pi_1 \rightarrow \cdots \rightarrow \pi_n \rightarrow \pi' \rightarrow 0.$$

We let  $D(G)$  denote the derived category of  $\text{Mod}_G^{\text{adm}}(R)$  ([Nee90, Kel96, Böh10]). The results of [Ver96, § III.3.2] on the Yoneda construction carry over to this setting (see e.g. [Pos11, Proposition A.13]), hence a natural  $R$ -linear isomorphism

$$\text{Ext}_G^n(\pi', \pi) \cong \text{Hom}_{D(G)}(\pi', \pi[n]).$$

*Proof of Corollary 2.* Since  $\text{Ind}_P^G$  and  $\text{Ord}_P$  are exact adjoint functors between  $\text{Mod}_M^{\text{adm}}(R)$  and  $\text{Mod}_G^{\text{adm}}(R)$  by Theorem 1, they induce adjoint functors between  $D(M)$  and  $D(G)$ , hence natural  $R$ -linear isomorphisms

$$\begin{aligned} \text{Ext}_M^n(\sigma, \text{Ord}_P(\pi)) &\cong \text{Hom}_{D(M)}(\sigma, \text{Ord}_P(\pi)[n]) \\ &\cong \text{Hom}_{D(G)}(\text{Ind}_P^G(\sigma), \pi[n]) \\ &\cong \text{Ext}_G^n(\text{Ind}_P^G(\sigma), \pi) \end{aligned}$$

for all  $n \geq 0$ .  $\square$

*Remark.* We give a more explicit proof of Corollary 2. The exact functor  $\text{Ind}_P^G$  and the counit of the adjunction between  $\text{Ind}_P^G$  and  $\text{Ord}_P$  induce an  $R$ -linear morphism

$$(15) \quad \text{Ext}_M^n(\sigma, \text{Ord}_P(\pi)) \rightarrow \text{Ext}_G^n(\text{Ind}_P^G(\sigma), \pi).$$

In the other direction, the exact (by Theorem 1) functor  $\text{Ord}_P$  and the unit of the adjunction between  $\text{Ind}_P^G$  and  $\text{Ord}_P$  induce an  $R$ -linear morphism

$$(16) \quad \text{Ext}_G^n(\text{Ind}_P^G(\sigma), \pi) \rightarrow \text{Ext}_M^n(\sigma, \text{Ord}_P(\pi)).$$

We prove that (16) is the inverse of (15). For  $n = 0$  this is the unit-counit equations. Assume  $n \geq 1$  and let

$$(17) \quad 0 \rightarrow \text{Ord}_P(\pi) \rightarrow \sigma_1 \rightarrow \cdots \rightarrow \sigma_n \rightarrow \sigma \rightarrow 0$$

be an exact sequence of admissible  $R$ -representations of  $M$ . By [Yon60, § 3], the image of the class of (17) under (15) is the class of any exact sequence of admissible  $R$ -representations of  $G$

$$(18) \quad 0 \rightarrow \pi \rightarrow \pi_1 \rightarrow \cdots \rightarrow \pi_n \rightarrow \text{Ind}_P^G(\sigma) \rightarrow 0$$

such that there exists a commutative diagram of admissible  $R$ -representations of  $G$

$$\begin{array}{ccccccccccc} 0 & \rightarrow & \text{Ind}_P^G(\text{Ord}_P(\pi)) & \rightarrow & \text{Ind}_P^G(\sigma_1) & \rightarrow & \cdots & \rightarrow & \text{Ind}_P^G(\sigma_n) & \rightarrow & \text{Ind}_P^G(\sigma) & \rightarrow & 0 \\ & & \downarrow & & \downarrow & & & & \downarrow & & \parallel & & \\ 0 & \longrightarrow & \pi & \longrightarrow & \pi_1 & \longrightarrow & \cdots & \longrightarrow & \pi_n & \longrightarrow & \text{Ind}_P^G(\sigma) & \longrightarrow & 0 \end{array}$$

in which the upper row is obtained from (17) by applying the exact functor  $\text{Ind}_P^G$ , the lower row is (18), and the leftmost vertical arrow is the natural morphism induced by the counit of the adjunction between  $\text{Ind}_P^G$  and  $\text{Ord}_P$ . Applying the exact functor  $\text{Ord}_P$  to the diagram and using the unit of the adjunction between  $\text{Ind}_P^G$  and  $\text{Ord}_P$  yields a commutative diagram of admissible  $R$ -representations of  $M$

$$\begin{array}{ccccccccccc} 0 & \rightarrow & \text{Ord}_P(\pi) & \longrightarrow & \sigma_1 & \longrightarrow & \cdots & \longrightarrow & \sigma_n & \longrightarrow & \sigma & \longrightarrow & 0 \\ & & \parallel & & \downarrow & & & & \downarrow & & \downarrow & & \\ 0 & \rightarrow & \text{Ord}_P(\pi) & \rightarrow & \text{Ord}_P(\pi_1) & \rightarrow & \cdots & \rightarrow & \text{Ord}_P(\pi_n) & \rightarrow & \text{Ord}_P(\text{Ind}_P^G(\sigma)) & \rightarrow & 0 \end{array}$$

in which the lower row is obtained from (18) by applying the exact functor  $\text{Ord}_P$ , the upper row is (17), and the rightmost vertical arrow is the natural morphism induced by the unit of the adjunction between  $\text{Ind}_P^G$  and  $\text{Ord}_P$ . The leftmost vertical morphism is the identity by the unit-counit equations. Thus the image of the class of (18) under (16) is the class of (17) by [Yon60, § 3]. We have proved that (16) is a left inverse of (15). The proof that it is a right inverse is dual.

**Corollary 5.** *Assume  $R$  artinian,  $p$  nilpotent in  $R$ , and  $\text{char}(F) = p$ . Let  $\sigma$  and  $\sigma'$  be two admissible  $R$ -representations of  $M$ . The functor  $\text{Ind}_P^G$  induces an  $R$ -linear isomorphism*

$$\text{Ext}_M^n(\sigma', \sigma) \xrightarrow{\sim} \text{Ext}_G^n(\text{Ind}_P^G(\sigma'), \text{Ind}_P^G(\sigma))$$

for all  $n \geq 0$ .

*Proof.* The isomorphism in the statement is the composite

$$\text{Ext}_M^n(\sigma', \sigma) \xrightarrow{\sim} \text{Ext}_M^n(\sigma', \text{Ord}_P(\text{Ind}_P^G(\sigma))) \xrightarrow{\sim} \text{Ext}_G^n(\text{Ind}_P^G(\sigma'), \text{Ind}_P^G(\sigma))$$

where the first isomorphism is induced by the unit of the adjunction between  $\text{Ind}_P^G$  and  $\text{Ord}_P$ , which is an isomorphism, and the second one is the isomorphism of Corollary 2 with  $\sigma'$  and  $\text{Ind}_P^G(\sigma)$  instead of  $\sigma$  and  $\pi$  respectively.  $\square$

We fix a minimal parabolic subgroup  $\mathbf{B} \subseteq \mathbf{G}$ , a maximal split torus  $\mathbf{S} \subseteq \mathbf{B}$ , and we write  $\Delta$  for the set of simple roots of  $\mathbf{S}$  in  $\mathbf{B}$ . We say that a parabolic subgroup  $\mathbf{P} = \mathbf{M}\mathbf{N}$  of  $\mathbf{G}$  is *standard* if  $\mathbf{B} \subseteq \mathbf{P}$  and  $\mathbf{S} \subseteq \mathbf{M}$ . In this case, we write  $\Delta_{\mathbf{P}}$  for the corresponding subset of  $\Delta$ , and given  $\alpha \in \Delta_{\mathbf{P}}$  (resp.  $\alpha \in \Delta \setminus \Delta_{\mathbf{P}}$ ) we write  $\mathbf{P}^{\alpha} = \mathbf{M}^{\alpha}\mathbf{N}^{\alpha}$  (resp.  $\mathbf{P}_{\alpha} = \mathbf{M}_{\alpha}\mathbf{N}_{\alpha}$ ) for the standard parabolic subgroup corresponding to  $\Delta_{\mathbf{P}} \setminus \{\alpha\}$  (resp.  $\Delta_{\mathbf{P}} \sqcup \{\alpha\}$ ).

Let  $C$  be an algebraically closed field of characteristic  $p$ . Given a standard parabolic subgroup  $P = MN$  and a smooth  $C$ -representation  $\sigma$  of  $M$ , there exists a largest standard parabolic subgroup  $P(\sigma) = M(\sigma)N(\sigma)$  such that the inflation of  $\sigma$  to  $P$  extends to a smooth  $C$ -representation  ${}^e\sigma$  of  $P(\sigma)$ , and this extension is unique ([AHHV17, II.7 Corollary 1]). We say that a smooth  $C$ -representation of  $G$  is *supercuspidal* if it is irreducible, admissible, and does not appear as a subquotient of  $\text{Ind}_P^G(\sigma)$  for any proper parabolic subgroup  $P = MN$  of  $G$  and any irreducible admissible  $C$ -representation  $\sigma$  of  $M$ . A *supercuspidal standard  $C[G]$ -triple* is a triple  $(P, \sigma, Q)$  where  $P = MN$  is a standard parabolic subgroup,  $\sigma$  is a supercuspidal  $C$ -representation of  $M$ , and  $Q$  is a parabolic subgroup of  $G$  such that  $P \subseteq Q \subseteq P(\sigma)$ . To such a triple is attached in [AHHV17] a smooth  $C$ -representation of  $G$

$$\mathbf{I}_G(P, \sigma, Q) := \text{Ind}_{P(\sigma)}^G({}^e\sigma \otimes \text{St}_Q^{P(\sigma)})$$

where  $\text{St}_Q^{P(\sigma)} := \text{Ind}_Q^{P(\sigma)}(1) / \sum_{Q \subsetneq Q' \subseteq P(\sigma)} \text{Ind}_{Q'}^{P(\sigma)}(1)$  (here 1 denotes the trivial  $C$ -representation) is the inflation to  $P(\sigma)$  of the generalised Steinberg representation of  $M(\sigma)$  with respect to  $M(\sigma) \cap Q$  ([GK14, Ly15]). It is irreducible and admissible ([AHHV17, I.3 Theorem 1]).

**Proposition 6.** *Assume  $\text{char}(F) = p$ . Let  $(P, \sigma, Q)$  and  $(P', \sigma', Q')$  be two supercuspidal standard  $C[G]$ -triples. If  $Q \not\subseteq Q'$ , then the  $C$ -vector space*

$$\text{Ext}_G^1(\mathbf{I}_G(P', \sigma', Q'), \mathbf{I}_G(P, \sigma, Q))$$

*is non-zero if and only if  $P' = P$ ,  $\sigma' \cong \sigma$ , and  $Q' = Q^{\alpha}$  for some  $\alpha \in \Delta_Q$ , in which case it is one-dimensional and the unique (up to isomorphism) non-split extension of  $\mathbf{I}_G(P', \sigma', Q')$  by  $\mathbf{I}_G(P, \sigma, Q)$  is the admissible  $C$ -representation of  $G$*

$$\text{Ind}_{P(\sigma)^{\alpha}}^G(\mathbf{I}_{M(\sigma)^{\alpha}}(M(\sigma)^{\alpha} \cap P, \sigma, M(\sigma)^{\alpha} \cap Q)).$$

*Proof.* There is a natural short exact sequence of admissible  $C$ -representations of  $G$

$$(19) \quad 0 \rightarrow \sum_{Q' \subsetneq Q'' \subseteq P(\sigma')} \text{Ind}_{Q''}^G(\sigma') \rightarrow \text{Ind}_{Q'}^G(\sigma') \rightarrow \mathbf{I}_G(P', \sigma', Q') \rightarrow 0.$$

Note that we can restrict the sum to those  $Q''$  that are minimal, i.e. of the form  $Q'_{\alpha}$  for some  $\alpha \in \Delta_{P(\sigma')} \setminus \Delta_{Q'}$ . Moreover, we deduce from [AHV17, Theorem 3.2] that its cosocle is isomorphic to  $\bigoplus_{\alpha \in \Delta_{P(\sigma')} \setminus \Delta_{Q'}} \mathbf{I}_G(P', \sigma', Q'_{\alpha})$ . Now if  $Q \not\subseteq Q'$ , then  $\text{Ord}_{\bar{Q}}(\mathbf{I}_G(P, \sigma, Q)) = 0$  by [AHV17, Theorem 1.1 (ii) and Corollary 4.13] so that using Corollary 2, we see that the long exact sequence of Yoneda extensions obtained by applying the functor  $\text{Hom}_G(-, \mathbf{I}_G(P, \sigma, Q))$  to (19) yields a natural  $C$ -linear isomorphism

$$\begin{aligned} \text{Ext}_G^{n-1}(\sum_{Q' \subsetneq Q'' \subseteq P(\sigma')} \text{Ind}_{Q''}^G(\sigma'), \mathbf{I}_G(P, \sigma, Q)) \\ \xrightarrow{\sim} \text{Ext}_G^n(\mathbf{I}_G(P', \sigma', Q'), \mathbf{I}_G(P, \sigma, Q)) \end{aligned}$$

for all  $n \geq 1$ . In particular, with  $n = 1$  and using the identification of the cosocle of the sum and [AHHV17, I.3 Theorem 2], we deduce that the  $C$ -vector space in the statement is non-zero if and only if  $P' = P$ ,  $\sigma' \cong \sigma$ , and  $Q = Q'_{\alpha}$  for some  $\alpha \in \Delta_{P(\sigma')} \setminus \Delta_{Q'}$  (or equivalently  $Q' = Q^{\alpha}$  for some  $\alpha \in \Delta_Q$ ), in which case it is one-dimensional. Finally, using again [AHV17, Theorem 3.2], we see that for all  $\alpha \in \Delta_Q$  the admissible  $C$ -representation of  $G$  in the statement is a non-split extension of  $\mathbf{I}_G(P, \sigma, Q^{\alpha})$  by  $\mathbf{I}_G(P, \sigma, Q)$ .  $\square$

**Corollary 7.** *Assume  $\text{char}(F) = p$ . Let  $\pi$  and  $\pi'$  be two irreducible admissible  $C$ -representations of  $G$ . If  $\pi$  is supercuspidal and  $\pi'$  is not the extension to  $G$  of a supercuspidal representation of a Levi subgroup of  $G$ , then  $\text{Ext}_G^1(\pi', \pi) = 0$ .*

*Proof.* By [AHHV17, I.3 Theorem 3], there exist two supercuspidal standard  $C[G]$ -triples  $(P, \sigma, Q)$  and  $(P', \sigma', Q')$  such that  $\pi \cong I_G(P, \sigma, Q)$  and  $\pi' \cong I_G(P', \sigma', Q')$ . The assumptions on  $\pi$  and  $\pi'$  are equivalent to  $P = G$  and  $Q' \neq G$ . In particular,  $Q \not\subseteq Q'$  and  $P \neq P'$  so that  $\text{Ext}_G^1(\pi', \pi) = 0$  by Proposition 6.  $\square$

## REFERENCES

- [AHHV17] N. Abe, G. Henniart, F. Herzig, and M.-F. Vignéras, *A classification of irreducible admissible mod  $p$  representations of  $p$ -adic reductive groups*, J. Amer. Math. Soc. **30** (2017), no. 2, 495–559.
- [AHV17] N. Abe, G. Henniart, and M.-F. Vignéras, *Modulo  $p$  representations of reductive  $p$ -adic groups: functorial properties*, e-print arXiv:1703.05599v2, to appear in Transactions of the American Mathematical Society, 2017.
- [Ber87] J. Bernstein, *Second adjointness for representations of reductive  $p$ -adic groups*, unpublished, available at <http://www.math.tau.ac.il/~bernstei>, 1987.
- [Büh10] T. Bühler, *Exact categories*, Expo. Math. **28** (2010), no. 1, 1–69.
- [CGP15] B. Conrad, O. Gabber, and G. Prasad, *Pseudo-reductive Groups*, second ed., New Mathematical Monographs, vol. 17, Cambridge University Press, Cambridge, 2015.
- [Dat09] J.-F. Dat, *Finitude pour les représentations lisses de groupes  $p$ -adiques*, J. Inst. Math. Jussieu **8** (2009), no. 2, 261–333.
- [Eme10a] M. Emerton, *Ordinary parts of admissible representations of  $p$ -adic reductive groups I. Definition and first properties*, Astérisque **331** (2010), 355–402.
- [Eme10b] ———, *Ordinary parts of admissible representations of  $p$ -adic reductive groups II. Derived functors*, Astérisque **331** (2010), 403–459.
- [GK14] E. Grosse-Klönne, *On special representations of  $p$ -adic reductive groups*, Duke Math. J. **163** (2014), no. 12, 2179–2216.
- [Hau16a] J. Hauseux, *Extensions entre séries principales  $p$ -adiques et modulo  $p$  de  $G(F)$* , J. Inst. Math. Jussieu **15** (2016), no. 2, 225–270.
- [Hau16b] ———, *Parabolic induction and extensions*, e-print arXiv:1607.02031, to appear in Algebra & Number Theory, 2016.
- [Hau16c] ———, *Sur une conjecture de Breuil-Herzig*, to appear in Journal für die reine und angewandte Mathematik, published online doi:10.1515/crelle-2016-0039, 2016.
- [Hu12] Y. Hu, *Diagrammes canoniques et représentations modulo  $p$  de  $\text{GL}_2(F)$* , J. Inst. Math. Jussieu **11** (2012), no. 1, 67–118.
- [Hu17] ———, *An application of a theorem of Emerton to mod  $p$  representations of  $\text{GL}_2$* , J. London Math. Soc. (2) **96** (2017), no. 3, 545–564.
- [Kel96] B. Keller, *Derived categories and their uses*, Handbook of Algebra, Volume 1, Elsevier, Amsterdam, 1996, pp. 671–701.
- [Ly15] T. Ly, *Représentations de Steinberg modulo  $p$  pour un groupe réductif sur un corps local*, Pacific J. Math. **277** (2015), no. 2, 425–462.
- [Nee90] A. Neeman, *The derived category of an exact category*, J. Algebra **135** (1990), no. 2, 388–394.
- [NSW08] J. Neukirch, A. Schmidt, and K. Wingberg, *Cohomology of Number Fields*, second ed., Grundlehren der Mathematischen Wissenschaften, vol. 323, Springer, Berlin, 2008.
- [Pos11] L. Positselski, *Mixed Artin-Tate motives with finite coefficients*, Moscow Math. J. **11** (2011), no. 2, 317–402.
- [Qui73] D. Quillen, *Higher algebraic K-theory: I*, Algebraic K-Theory I, Lecture Notes in Mathematics, vol. 341, Springer, Berlin, 1973, pp. 85–147.
- [Ver96] J.-L. Verdier, *Des catégories dérivées des catégories abéliennes*, Astérisque **239** (1996).
- [Vig96] M.-F. Vignéras, *Représentations  $l$ -modulaires d’un groupe réductif  $p$ -adique avec  $l \neq p$* , Progress in Mathematics, vol. 137, Birkhäuser, Boston, 1996.
- [Vig16] M.-F. Vignéras, *The right adjoint of the parabolic induction*, Arbeitstagung Bonn 2013, Progress in Mathematics, vol. 319, Birkhäuser, Basel, 2016, pp. 405–425.
- [Yon60] N. Yoneda, *On Ext and exact sequences*, J. Fac. Sci. Univ. Tokyo Sect. 1A Math. **8** (1960), 507–576.