



Yet another Hopf invariant

Jean-Paul Doeraene¹ Mohammed El Haouari¹

Received: April 20, 2016/Accepted: September 10, 2018/Online: December 2, 2018

Abstract

The classical Hopf invariant is defined for a map $f : S^r \rightarrow X$. Here we define ‘hcat’ which is some kind of Hopf invariant built with a construction in Ganea’s style, valid for maps not only on spheres but more generally on a ‘relative suspension’ $f : \Sigma_A W \rightarrow X$. We study the relation between this invariant and the sectional category and the relative category of a map. In particular, for $\iota_X : A \rightarrow X$ being the ‘restriction’ of f on A , we have $\text{relcat } \iota_X \leq \text{hcat } f \leq \text{relcat } \iota_X + 1$ and $\text{relcat } f \leq \text{hcat } f$.

Keywords: Ganea fibration, sectional category, Hopf invariant..

msc: 55M30.

Our aim here is to make clearer the link between the Lusternik-Schnirelmann category (cat), more generally the ‘relative category’ (relcat), closely related to James’ sectional category (secat), and the Hopf invariants. In order to do this, we introduce a new integer, namely hcat , that combines the Iwaze’s version of Hopf invariant², based on the *difference up to homotopy between two maps* defined for a given section of a Ganea fibration, and the framework of the sectional and relative categories, searching for the *least integer* such that the Ganea fibration has a section, possibly with additional conditions. To do this combination, we simply define our invariant hcat , as the least integer such that the Ganea fibration has a section σ with additional condition that the corresponding two maps ($f \circ \sigma$ and ω_n in this paper) are homotopic.

It appears that for $f : S^r \rightarrow X$ or even for $f : \Sigma W \rightarrow X$, we obtain an integer that can be either $\text{cat}(X)$, or $\text{cat}(X) + 1$. More generally, for any $f : \Sigma_A W \rightarrow X$, we have $\text{relcat}(f \circ \theta) \leq \text{hcat}(f) \leq \text{relcat}(f \circ \theta) + 1$, where $\theta : A \rightarrow \Sigma_A W$ is the map arising in the construction of $\Sigma_A W$.

In section 2, we study the influence of hcat in a homotopy pushout. In section 3, we introduce the ‘strong’ version of our invariant, and we obtain another important inequality: for any $f : \Sigma_A W \rightarrow X$, we have $\text{relcat}(f) \leq \text{hcat}(f)$. In section 4, we give alternative equivalent conditions to get hcat . Applications and examples are given.

¹Département de Mathématiques, UMR-CNRS 8524, Université de Lille, 59655 Villeneuve d’Ascq Cedex, France


²Iwase, 1198, “Ganea’s conjecture on Lusternik-Schnirelmann category”.

1 The Hopf category

We work in the category of pointed topological spaces. All constructions are made up to homotopy. A ‘homotopy commutative diagram’ has to be understood in the sense of Mather.

Recall the following construction:

Definition 1 – For any map $\iota_X : A \rightarrow X$, the *Ganea construction* of ι_X is the following sequence of homotopy commutative diagrams ($i \geq 0$):



where the outside square is a homotopy pullback, the inside square is a homotopy pushout and the map $g_{i+1} = (g_i, \iota_X) : G_{i+1} \rightarrow X$ is the whisker map induced by this homotopy pushout. The iteration starts with $g_0 = \iota_X : A \rightarrow X$. We set $\alpha_0 = \text{id}_A$.

For any $i \geq 0$, there is a whisker map $\theta_i = (\text{id}_A, \alpha_i) : A \rightarrow F_i$ induced by the homotopy pullback. Thus we have the sequence of maps $A \xrightarrow{\theta_i} F_i \xrightarrow{\eta_i} A$ and θ_i is a homotopy section of η_i . Moreover we have $\gamma_i \circ \alpha_i \simeq \alpha_{i+1}$, thus also $\alpha_{i+1} \simeq \gamma_i \circ \gamma_{i-1} \circ \dots \circ \gamma_0$.

We denote by $\gamma_{i,j} : G_i \rightarrow G_j$ the composite $\gamma_{j-1} \circ \dots \circ \gamma_{i+1} \circ \gamma_i$ (for $i < j$) and set $\gamma_{i,i} = \text{id}_{G_i}$.

Of course, everything in the Ganea construction depends on ι_X . We sometimes denote G_i by $G_i(\iota_X)$ to avoid ambiguity.

Definition 2 – Let $\iota_X : A \rightarrow X$ be any map.

1) The *sectional category* of ι_X is the least integer n such that the map $g_n : G_n(\iota_X) \rightarrow X$ has a homotopy section, i.e. there exists a map $\sigma : X \rightarrow G_n(\iota_X)$ such that $g_n \circ \sigma \simeq \text{id}_X$.

2) The *relative category* of ι_X is the least integer n such that the map $g_n : G_n(\iota_X) \rightarrow X$ has a homotopy section σ and $\sigma \circ \iota_X \simeq \alpha_n$.

3) The *relative category of order k* of ι_X is the least integer n such that the map $g_n : G_n(\iota_X) \rightarrow X$ has a homotopy section σ and $\sigma \circ g_k \simeq \gamma_{k,n}$.

We denote the sectional category by $\text{secat}(\iota_X)$, the relative category by $\text{relcat}(\iota_X)$, and the relative category of order k by $\text{relcat}_k(\iota_X)$. If $A = *$, $\text{secat}(\iota_X) = \text{relcat}(\iota_X)$

1. The Hopf category

and is denoted simply by $\text{cat}(X)$; this is the ‘normalized’ version of the Lusternik-Schnirelmann category.

Clearly, $\text{secat}(\iota_X) \leq \text{relcat}(\iota_X)$. We have also $\text{relcat}(\iota_X) \leq \text{relcat}_1(\iota_X)$, see Proposition 1 below.

In the sequel, we will consider a given homotopy pushout:

$$\begin{array}{ccc} W & \xrightarrow{\eta} & A \\ \beta \downarrow & & \downarrow \theta \\ A & \xrightarrow{\theta} & \Sigma_A W \end{array}$$

In other words, the map θ is a map such that $\text{Pushcat } \theta \leq 1$ in the sense of Doeraene and El Haouari. We call this homotopy pushout a ‘relative suspension’ because in some sense, A plays the role of the point in the ordinary suspension.

We also consider any map $f: \Sigma_A W \rightarrow X$, and set $\iota_X = f \circ \theta$.

We don’t assume $\eta \simeq \beta$ in general. This is true, however, if θ is a homotopy monomorphism, and in this case we can ‘think’ of ι_X as the ‘restriction’ of f on A .

For $n \geq 1$, consider the following homotopy commutative diagram:

$$\begin{array}{ccccc} & & W & \xrightarrow{\beta} & A \\ & \swarrow \eta & & & \swarrow \theta \\ A & \xrightarrow{\theta} & \Sigma_A W & \xrightarrow{\theta} & \Sigma_A W \\ & \downarrow \theta & \downarrow \omega_n & \downarrow \alpha_{n-1} & \downarrow f \\ & F_{n-1}(\iota_X) & \xrightarrow{\omega_n} & G_{n-1}(\iota_X) & \downarrow g_{n-1} \\ & \swarrow \alpha_n & \downarrow \gamma_{n-1} & \swarrow g_n & X \\ A & \xrightarrow{\alpha_n} & G_n(\iota_X) & \xrightarrow{g_n} & X \end{array} \quad (\dagger)$$

where the map $W \rightarrow F_{n-1}$ is induced by the bottom outer homotopy pullback and the map $\omega_n: \Sigma_A W \rightarrow G_n$ is induced by the top inner homotopy pushout. We have $f \simeq g_n \circ \omega_n$ by the ‘Whisker maps inside a cube’ lemma (see Doeraene and El Haouari (2013), Lemma 49). Also notice that $\alpha_n \simeq \omega_n \circ \theta \simeq \gamma_{n-1} \circ \alpha_{n-1}$; so $\omega_n \simeq (\alpha_n, \alpha_n)$ is the whisker map of two copies of α_n induced by the homotopy pushout $\Sigma_A W$. Finally, for all $k \geq 1$, we can see that $\omega_n \simeq \gamma_{k,n} \circ \omega_k$.

Definition 3 – The Hopf category of f is the least integer $n \geq 1$ such that $g_n: G_n(\iota_X) \rightarrow X$ has a homotopy section $\sigma: X \rightarrow G_n(\iota_X)$ such that $\sigma \circ f \simeq \omega_n$.

We denote this integer by $\text{hcat}(f)$.

Actually, speaking of ‘Hopf category of f ’ is a misuse of language. We should speak of ‘Hopf category of the datas η, β and f ’.

Example 1 – Let $X = \Sigma_A W$ and $f \simeq \text{id}_X$. Then, as might be expected, $\text{hcat}(f) = 1$. Indeed, in this case, as $g_1 \circ \omega_1 \simeq f \simeq \text{id}_X$, ω_1 is a homotopy section of g_1 . Moreover, $\omega_1 \circ f \simeq \omega_1 \circ \text{id}_X \simeq \omega_1$, so $\text{hcat}(f) = 1$.

Example 2 – Let $X \neq *$ and $W = A \vee A$, $\beta \simeq \text{pr}_1 : A \vee A \rightarrow A$ and $\eta \simeq \text{pr}_2 : A \vee A \rightarrow A$ the obvious maps. Then $\Sigma_A W \simeq *$ and we have no choice for f that must be the null map $f : * \rightarrow X$. In this case the condition $\sigma \circ f \simeq \omega_n$ is always satisfied, so $\text{hcat}(f) = \text{secat}(\iota_X) = \text{cat}(X)$.


Notice that relcat is a particular case of hcat : When $W = A$, $\eta \simeq \beta \simeq \text{id}_A$, then $\iota_X \simeq f$, $\omega_n \simeq \alpha_n$ and $\text{hcat}(f) = \text{relcat}(\iota_X)$. Also relcat_1 is a particular case of hcat : When $W = F_0$, then $\Sigma_A W \simeq G_1$, $\theta \simeq \gamma_0 \simeq \alpha_1$, and if, moreover, $f \simeq g_1$, then $\omega_n \simeq \gamma_{1,n}$ and $\text{hcat}(f) = \text{relcat}_1(\iota_X)$.

The following proposition shows that these particular cases are in fact lower and upper bounds for $\text{hcat}(f)$.

Proposition 1 – *Whatever can be f (and $\iota_X = f \circ \theta$), we have*


$$\text{secat}(f) \leq \text{relcat}(\iota_X) \leq \text{hcat}(f) \leq \text{relcat}_1(\iota_X) \leq \text{relcat}(\iota_X) + 1.$$

Proof. Consider the following homotopy commutative diagram ($n \geq 1$):



We see that if there is a map $\sigma : X \rightarrow G_n$ such that $\omega_n \simeq \sigma \circ f$ then $\alpha_n \simeq \sigma \circ \iota_X$ and this proves the second inequality.

Now consider the following homotopy commutative diagram ($n \geq 1$):



We see that if there is a map $\sigma : X \rightarrow G_n$ such that $\gamma_{1,n} \simeq \sigma \circ g_1$ then $\omega_n \simeq \sigma \circ f$ and this proves the third inequality.

The first inequality comes from $\text{secat}(f) \leq \text{secat}(\iota_X) \leq \text{relcat}(\iota_X)$, the first of these two inequalities comes from Doeraene and El Haouari (2013), Proposition 29.

Finally, the fourth inequality is proved in Doeraene (2016). □

2. Hopf invariant and homotopy pushout

So $\text{hcat}(f)$ establishes a ‘dichotomy’ between maps $f: \Sigma_A W \rightarrow X$:

- Either $\text{hcat}(f) = \text{relcat}(t_X)$ and we have a σ such that $f \circ \sigma \simeq \omega_n$ already for $n = \text{secat}(t_X)$;
- either $\text{hcat}(f) = \text{relcat}(t_X) + 1$ and we have a σ such that $f \circ \sigma \simeq \omega_n$ only for $n > \text{secat}(t_X)$

Our last example of the section shows that the inequalities of Proposition 1 can be strict, and even that two may be strict at the same time:

Example 3 – Let $X = *$, $A \neq *$ and consider $t_*: A \rightarrow *$. We have $G_i(t_*) \simeq A \bowtie \dots \bowtie A$, the join of $i + 1$ copies of A . For any k , $\gamma_{k,k} \simeq \text{id}$, so it cannot factorize through $*$; but $\gamma_{k,k+1}$ is homotopic to the null map, so $\text{relcat}_k(t_*) = k + 1$. Now consider $f \simeq g_1(t_*): A \bowtie A \rightarrow *$. As said before, in this case we have $\text{hcat}(f) = \text{relcat}_1(t_X)$. So we get $\text{secat}(f) = 0 < \text{relcat}(t_*) = 1 < \text{hcat}(f) = \text{relcat}_1(t_*) = 2$.

2 Hopf invariant and homotopy pushout

Let us consider any homotopy commutative square:


$$\begin{array}{ccc} \Sigma_A W & \xrightarrow{\rho} & B \\ f \downarrow & & \downarrow \kappa_Y \\ X & \xrightarrow{\chi} & Y \end{array} \quad (\ddagger)$$

Proposition 2 – *The homotopy commutative square above can be splitted into the following homotopy commutative diagram:*


$$\begin{array}{ccccccc} & & & & f & & \\ & & & & \curvearrowright & & \\ \Sigma_A W & \longrightarrow & G_1(t_X) & \longrightarrow & G_n(t_X) & \longrightarrow & X \\ \downarrow \rho & & \downarrow & & \downarrow & & \downarrow \chi \\ B & \longrightarrow & G_1(\kappa_Y) & \longrightarrow & G_n(\kappa_Y) & \longrightarrow & Y \\ & & & & \curvearrowleft & & \\ & & & & \kappa_Y & & \end{array}$$

Proof. Set $\phi = \rho \circ \theta$. Since $\theta \circ \eta \simeq \theta \circ \beta$, also $\phi \circ \eta \simeq \phi \circ \beta$. First notice that we can insert the original homotopy square inside the following homotopy commutative

diagram:




By induction on $n \geq 1$, starting from the outside cube of the above diagram and $\phi_0 = \phi$, we can build a homotopy diagram:




where the dashed and dotted maps are induced by the homotopy pullback $F_{n-1}(\kappa_Y)$ and the homotopy pushout $G_n(t_X)$ respectively.

So we obtain a homotopy commutative diagram:



Finally take the homotopy pushout inside the upper and lower left squares to get the homotopy commutative diagram:



2. Hopf invariant and homotopy pushout

and this gives the required splitting of the original square. \square

Proposition 3 – *If the square \ddagger is a homotopy pushout, then*

$$\text{relcat}(\kappa_Y) \leq \text{hcat}(f).$$

As a particular case, when $B \simeq *$, Y is the homotopy cofibre of f , and $\text{relcat}(\kappa_Y) = \text{cat}(Y)$. So the Proposition asserts that $\text{hcat}(f) \geq \text{cat}(Y)$.

Proof. Let $\text{hcat}(f) \leq n$, so we have a homotopy section σ of $g_n(\iota_X)$ such that $\sigma \circ f \simeq \omega_n$. First apply the ‘Whisker maps inside a cube’ lemma to the outer part of the following homotopy commutative diagram:

$$\begin{array}{ccccc}
 & \Sigma_A W & \longrightarrow & B & \\
 \omega_n \swarrow & \parallel & & \parallel & \searrow \alpha_n \\
 G_n(\iota_X) & \longrightarrow & S & \xrightarrow{b} & G_n(\kappa_Y) \\
 \downarrow & \parallel & \downarrow c & \parallel & \downarrow g_n \\
 X & \longrightarrow & Y & \xrightarrow{=} & Y \\
 & \Sigma_A W & \longrightarrow & B & \\
 & \swarrow & & \swarrow & \\
 & X & & Y &
 \end{array}$$

where the inner horizontal squares are homotopy pushouts, and c and b are the whisker maps induced by the homotopy pushout S . Next build the following homotopy commutative diagram:

$$\begin{array}{ccccc}
 & \Sigma_A W & \longrightarrow & B & \\
 f \swarrow & \downarrow & & \downarrow \kappa & \\
 X & \longrightarrow & Y & & \\
 \downarrow \sigma & & \downarrow & & \\
 G_n(\iota_X) & \longrightarrow & S & \xrightarrow{b} & G_n(\kappa_Y) \\
 & \swarrow \omega_n & & \swarrow a & \searrow \alpha_n \\
 & \Sigma_A W & \longrightarrow & B & \\
 & \downarrow d & & \downarrow & \\
 & X & & Y &
 \end{array}$$

where d is the whisker map induced by the homotopy pushout Y . Let $\sigma' = b \circ d$. We have $g_n \circ \sigma' \simeq g_n \circ b \circ d \simeq c \circ d \simeq \text{id}_C$ and $\sigma' \circ \kappa_Y \simeq b \circ d \circ \kappa_Y \simeq b \circ a \simeq \alpha_n$. \square

Corollary 1 – *In the diagram \ddagger , if $\text{relcat}(\kappa_Y) = \text{relcat}(\iota_X) + 1$, then $\text{hcat}(f) = \text{relcat}(\iota_X) + 1$.*

Proof. By Proposition 3, the hypothesis implies that $\text{hcat}(f) \geq \text{relcat}(\iota_X) + 1$. But by Proposition 1, we have $\text{hcat}(f) \leq \text{relcat}(\iota_X) + 1$. So we have the equality. \square

It is now easy to exhibit examples of maps f with $\text{hcat}(f) = \text{relcat}(\iota_X) + 1$. Indeed there are plenty examples of homotopy pushouts where $\text{relcat}(\kappa_Y) = \text{relcat}(\iota_X) + 1$:

Example 4 – Let $A = B = *$ and $f: S^r \rightarrow S^n$ be any of the Hopf maps $S^3 \rightarrow S^2$, $S^7 \rightarrow S^4$ or $S^{15} \rightarrow S^8$. So here $\text{relcat}(\iota_X) = \text{cat}(S^n) = 1$. On the other hand it is well known that those maps have a homotopy cofibre S^n/S^r of category 2, so here $\text{relcat}(\kappa_Y) = \text{cat}(S^n/S^r) = 2$. By Corollary 1, we have $\text{hcat}(f) = 2$.

Example 5 – Let f be the map u in the homotopy cofibration

$$Z \bowtie Z \xrightarrow{u} \Sigma Z \vee \Sigma Z \xrightarrow{t_1} \Sigma Z \times \Sigma Z$$

where $Z \bowtie Z \simeq \Sigma(Z \wedge Z)$ is the join of two copies of Z and is also the suspension of the smash product of two copies of Z . Let $A = B = *$, $\Sigma Z \neq *$. We have $\text{relcat}(\iota_X) = \text{cat}(\Sigma Z \vee \Sigma Z) = 1$ and $\text{relcat}(\kappa_Y) = \text{cat}(\Sigma Z \times \Sigma Z) = 2$, so by Corollary 1 again, we have $\text{hcat}(u) = 2$.

Example 6 – For $i \geq 1$, let f be the map β_i in the Ganea construction:

$$\begin{array}{ccccc} A & \xrightarrow{\theta_i} & F_i & \xrightarrow{\eta_i} & A \\ & \searrow \alpha_i & \downarrow \beta_i & & \downarrow \alpha_{i+1} \\ & & G_i & \xrightarrow{\gamma_i} & G_{i+1} \end{array}$$

Actually F_i is a join over A of $i + 1$ copies of F_0 , and also a relative suspension $\Sigma_A W$ where W is a relative smash product. For any $i \leq \text{relcat}(\iota_X)$, we have $\text{relcat}(\alpha_i) = i$, see Doeraene and El Haouari (2013), Proposition 23. So by Corollary 1 again, if $i < \text{relcat}(\iota_X)$, we have $\text{hcat}(\beta_i) = \text{relcat}(\alpha_i) + 1 = i + 1$.

3 The Strong Hopf category

In Doeraene and El Haouari (2013), we introduced the strong version of relcat , namely Relcat . In this section, we introduce the strong version of hcat , namely Hcat . This gives an alternative way, sometimes usefull, to see if a map has a Hopf category less or equal to n . Also this will lead to a new inequality: $\text{hcat}(f) \geq \text{relcat}(f)$. Consequently, if $\text{relcat}(f) > \text{relcat}(\iota_X)$, then $\text{hcat}(f) = \text{relcat}(\iota_X) + 1$.

Definition 4 – The *strong Hopf category* of a map $f: \Sigma_A W \rightarrow X$ is the least integer $n \geq 1$ such that:

- there are maps $\iota_0: A \rightarrow X_0$ and a homotopy inverse $\lambda: X_0 \rightarrow A$, i.e. $\iota_0 \circ \lambda \simeq \text{id}_{X_0}$ and $\lambda \circ \iota_0 \simeq \text{id}_A$;

3. The Strong Hopf category

- for each $i, 0 \leq i < n$, there is a homotopy commutative cube:

$$\begin{array}{ccccc}
 & & W & \xrightarrow{\beta} & A \\
 & \nearrow \eta & \downarrow & & \swarrow & \downarrow \iota_i \\
 A & \xrightarrow{\quad} & \Sigma_A W & & & \\
 \parallel & & \downarrow & \xrightarrow{z_i} & \downarrow \zeta_{i+1} & \downarrow \\
 & \swarrow & Z_i & \xrightarrow{\quad} & X_i & \\
 & & \downarrow & & \swarrow \chi_i & \\
 A & \xrightarrow{\iota_{i+1}} & X_{i+1} & & &
 \end{array} \tag{h}$$

where the bottom square is a homotopy pushout.

- $X_n = X$ and $\zeta_n \simeq f$.

We denote this integer by $\text{Hcat}(f)$.

Notice that $\iota_{i+1} \simeq \zeta_{i+1} \circ \theta \simeq \chi_i \circ \iota_i$. In particular, this means that $\text{Pushcat}(\iota_i) \leq i$ in the sense of Doeraene and El Haouari, Definition 3.

For $0 \leq i \leq n$, define the sequence of maps $\xi_i: X_i \rightarrow X$ with the relation $\xi_i = \xi_{i+1} \circ \chi_i$ (when $i < n$), starting with $\xi_n = \text{id}_X$. We have $\xi_n \circ \iota_n \simeq \iota_X$ and $\xi_i \circ \iota_i = \xi_{i+1} \circ \chi_i \circ \iota_i \simeq \xi_{i+1} \circ \iota_{i+1} \simeq \iota_X$ by decreasing induction. Also $\iota_X \circ \lambda \simeq \xi_0 \circ \iota_0 \circ \lambda \simeq \xi_0$. Moreover, for $0 < i \leq n$ we have we have $\xi_i \circ \zeta_i \simeq f$ by the ‘Whisker maps inside a cube lemma’. So we have the following homotopy diagram:

$$\begin{array}{ccccccc}
 & & W & \xrightarrow{\eta} & A & & \\
 & \swarrow \beta & \downarrow & & \swarrow & \searrow \theta & \\
 A & \xrightarrow{\quad} & \Sigma_A W & \xrightarrow{\quad} & \Sigma_A W & & \\
 \downarrow \iota_i & & \downarrow & \xrightarrow{\zeta_{i+1}} & \downarrow & & \downarrow f \\
 & \swarrow & Z_i & \xrightarrow{\quad} & A & \xrightarrow{\iota_X} & \\
 & & \downarrow & & \swarrow \iota_{i+1} & \searrow & \\
 X_i & \xrightarrow{\chi_i} & X_{i+1} & \xrightarrow{\xi_{i+1}} & X & &
 \end{array}$$

We say that a map $g: B \rightarrow Y$ is ‘relatively dominated’ by a map $f: B \rightarrow X$ if there is a map $\varphi: X \rightarrow Y$ with a homotopy section $\sigma: Y \rightarrow X$ such that $\varphi \circ f \simeq g$ and $\sigma \circ g \simeq f$.

Proposition 4 – A map $g: \Sigma_A W \rightarrow Y$ has $\text{hcat}(g) \leq n$ iff g is relatively dominated by a map $f: \Sigma_A W \rightarrow X$ with $\text{Hcat}(f) \leq n$.

Proof. Consider the map $\omega_n: \Sigma_A W \rightarrow G_n(\iota_Y)$ as in diagram † and notice that $\text{Hcat}(\omega_n) \leq n$. If $\text{hcat}(f) \leq n$, then f is relatively dominated by ω_n .

For the reverse direction, by hypothesis, we have a map φ and a homotopy section σ such that $\varphi \circ f \simeq g$ and $\sigma \circ g \simeq f$; composing with θ , we have also $\varphi \circ \iota_X \simeq \iota_Y$ and $\sigma \circ \iota_Y \simeq \iota_X$. From the hypothesis $\text{Hcat}(f) \leq n$, we get a sequence of homotopy commutative diagrams, for $0 \leq i < n$, which gives the top part of the following diagram.

We show by induction that the map $\varphi \circ \xi_i: X_i \rightarrow Y$ factors through $g_i: G_i(\iota_Y) \rightarrow Y$ up to homotopy. This is true for $i = 0$ since we have $\xi_0 \simeq \iota_X \circ \lambda$, so $\varphi \circ \xi_0 \simeq \varphi \circ \iota_X \circ \lambda \simeq \iota_Y \circ \lambda = g_0 \circ \lambda$. Suppose now that we have a map $\lambda_i: X_i \rightarrow G_i(\iota_Y)$ such that $g_i \circ \lambda_i \simeq \varphi \circ \xi_i$. Then we construct a homotopy commutative diagram

$$\begin{array}{ccccc}
 & & Z_i & \xrightarrow{\quad} & A \\
 & \swarrow z_i & \vdots & \nearrow i_{i+1} & \parallel \\
 X_i & \xrightarrow{\quad} & X_{i+1} & \xrightarrow{\quad} & X \\
 \downarrow \lambda_i & & \downarrow \lambda_{i+1} & \nearrow \xi_{i+1} & \parallel \\
 & & F_i & \xrightarrow{\quad} & A \\
 & \swarrow & \vdots & \searrow \alpha_{i+1} & \parallel \\
 G_i(\iota_Y) & \xrightarrow{\quad} & G_{i+1}(\iota_Y) & \xrightarrow{\quad} & Y \\
 & & & \searrow g_{i+1} & \parallel \\
 & & & & \varphi
 \end{array}$$

where $Z_i \rightarrow F_i$ is the whisker map induced by the bottom homotopy pullback and $\lambda_{i+1}: X_{i+1} \rightarrow G_{i+1}(\iota_Y)$ is the whisker map induced by the top homotopy pushout. The composite $g_{i+1} \circ \lambda_{i+1}$ is homotopic to $\varphi \circ \xi_{i+1}$. Hence the inductive step is proven.

At the end of the induction, we have $g_n \circ \lambda_n \simeq \varphi \circ \xi_n = \varphi \circ \text{id}_X = \varphi$. As we have a homotopy section $\sigma: Y \rightarrow X_n = X$ of φ , we get a homotopy section $\lambda_n \circ \sigma$ of g_n . Moreover, we have $(\lambda_n \circ \sigma) \circ g \simeq \lambda_n \circ f \simeq \lambda_n \circ \zeta_n \simeq \omega_n$. \square

Example 7 – If we consider any relative suspension $\Sigma_A f: \Sigma_A W \rightarrow \Sigma_A Z$ (and in particular, of course, when $A = *$, any suspension $\Sigma f: \Sigma W \rightarrow \Sigma Z$), we have $\text{Hcat}(\Sigma_A f) = 1$. And so, any map g that is relatively dominated by a (relative) suspension has $\text{hcat}(g) = 1$.

In fact, by definition, a map g has $\text{Hcat}(g) = 1$ if and only if g is a (relative) suspension. There are maps for which the strong Hopf category is greater than the Hopf category: For instance, consider the null map $f: * \rightarrow X$ of Example 2; if X is a space with $\text{cat}(X) = 1$ that is not a suspension, then f cannot be a suspension, so $\text{Hcat}(f) > \text{hcat}(f) = 1$.

Proposition 5 – In the diagram \mathfrak{h} , we have

$$\text{Relcat}(\zeta_i) \leq i$$

As ω_i is a particular case of ζ_i , this implies $\text{Relcat}(\omega_i) \leq i$.

3. The Strong Hopf category

Proof. For $i > 0$, let build the following homotopy diagram where the three squares are homotopy pushouts:

$$\begin{array}{ccccc}
 & & \eta & & \\
 & & \curvearrowright & & \\
 W & \longrightarrow & Z_{i-1} & \longrightarrow & A_\theta \\
 \downarrow \beta & & \downarrow & & \downarrow \\
 A & \longrightarrow & C_{i-1} & \xrightarrow{z_{i-1}} & \Sigma_A W \\
 \downarrow \iota_{i-1} & & \downarrow c_{i-1} & & \downarrow \zeta_i \\
 & & X_{i-1} & \longrightarrow & X_i
 \end{array}$$

and where the map $c_{i-1} = (\iota_{i-1}, z_{i-1})$ is the whisker map induced by the homotopy pushout.

We have $\text{secat}(\iota_{i-1}) \leq \text{Pushcat}(\iota_{i-1}) \leq i - 1$ by Doeraene and El Haouari (2013), Theorem 18. So $\text{secat}(c_{i-1}) \leq i - 1$ by Doeraene and El Haouari (2013), Proposition 29. So $\text{Relcat}(c_{i-1}) \leq (i - 1) + 1 = i$ by Doeraene and El Haouari 2013, Theorem 18. And this implies $\text{Relcat}(\zeta_i) \leq i$ by Doeraene and El Haouari (2013), Lemma 11. \square

Theorem 1 – For any $f: \Sigma_A W \rightarrow X$, we have

$$\text{Relcat}(f) \leq \text{Hcat}(f) \quad \text{and} \quad \text{relcat}(f) \leq \text{hcat}(f)$$

Proof. If $\text{Hcat}(f) = n$, then we have $f \simeq \zeta_n$ in \mathfrak{h} . So $\text{Relcat}(f) = \text{Relcat}(\zeta_n) \leq n$ by Proposition 5.

If $\text{hcat}(f) = n$, then f is relatively dominated by ω_n . As $\text{Relcat}(\omega_n) \leq n$, we have $\text{relcat}(f) \leq n$ by Doeraene and El Haouari (2013), Proposition 10. \square

As a corollary, we get an indirect proof of Proposition 3 because $\text{relcat}(\kappa_Y) \leq \text{relcat}(f)$ by Doeraene and El Haouari (2013), Lemma 11, that asserts that a homotopy pushout doesn't increase the relative category.

It is not difficult to find an example where these inequalities are strict:

Example 8 – Let f be the map t_1 in the homotopy cofibration

$$Z \bowtie Z \xrightarrow{u} \Sigma Z \vee \Sigma Z \xrightarrow{t_1} \Sigma Z \times \Sigma Z$$

Let $A = *$, $\Sigma Z \neq *$. As t_1 is a homotopy cofibre, we have $\text{relcat}(t_1) \leq \text{Relcat}(t_1) \leq 1$, see Doeraene and El Haouari (2013), Proposition 9. On the other hand, we have $\text{Hcat}(t_1) \geq \text{hcat}(t_1) \geq \text{relcat}(t_X) = \text{cat}(\Sigma Z \times \Sigma Z) = 2$ by Proposition 1.

4 Equivalent conditions to get the Hopf category

Let be given any map $f: \Sigma_A W \rightarrow X$ with $\text{secat}(\iota_X) \leq n$ and any homotopy section $\sigma: X \rightarrow G_n$ of $g_n: G_n \rightarrow X$. Consider the following homotopy pullbacks:

$$\begin{array}{ccccc}
 Q & \xrightarrow{\pi} & \Sigma_A W & & \\
 \downarrow \pi' & & \downarrow \theta_n^W & \searrow & \\
 \Sigma_A W & \xrightarrow{\bar{\sigma}} & H_n & \xrightarrow{\eta_n^W} & \Sigma_A W \\
 \downarrow f & & \downarrow f_n & & \downarrow f \\
 X & \xrightarrow{\sigma} & G_n & \xrightarrow{g_n} & X
 \end{array}$$

where $\theta_n^W = (\omega_n, \text{id}_{\Sigma_A X})$ is the whisker map induced by the homotopy pullback H_n . By the ‘Prism lemma’ (see Doeraene and El Haouari (2013), Lemma 46, for instance), we know that the homotopy pullback of σ and f_n is indeed $\Sigma_A W$, and that $\eta_n^W \circ \bar{\sigma} \simeq \text{id}_{\Sigma_A W}$. Also notice that $\pi \simeq \pi'$ since $\pi \simeq \eta_n^W \circ \theta_n^W \circ \pi \simeq \eta_n^W \circ \bar{\sigma} \circ \pi' \simeq \pi'$.

Proposition 6 – *Let be given any map $f: \Sigma_A W \rightarrow X$ with $\text{secat}(\iota_X) \leq n$ and any homotopy section $\sigma: X \rightarrow G_n(\iota_X)$ of $g_n: G_n(\iota_X) \rightarrow X$. With the same definitions and notations as above, the following conditions are equivalent:*

- (i) $\sigma \circ f \simeq \omega_n$.
- (ii) π has a homotopy section.
- (iii) π is a homotopy epimorphism.
- (iv) $\theta_n^W \simeq \bar{\sigma}$.

Proof. We have the following sequence of implications:

(i) \implies (ii): Since $\sigma \circ f \simeq \omega_n \simeq f_n \circ \theta_n^W \circ \text{id}_{\Sigma_A W}$, we have a whisker map $(f, \text{id}_{\Sigma_A W}): \Sigma_A W \rightarrow Q$ induced by the homotopy pullback Q which is a homotopy section of π .

(ii) \implies (iii): Obvious.

(iii) \implies (iv): We have $\theta_n^W \circ \pi \simeq \bar{\sigma} \circ \pi$ since $\pi \simeq \pi'$. Thus $\theta_n^W \simeq \bar{\sigma}$ since π is a homotopy epimorphism.

(iv) \implies (i): We have $\sigma \circ f \simeq f_n \circ \bar{\sigma} \simeq f_n \circ \theta_n^W \simeq \omega_n$. □

Theorem 2 – *Let be a $(q-1)$ -connected map $\iota_X: A \rightarrow X$ with $\text{secat} \iota_X \leq n$. If $\Sigma_A W$ is a CW-complex with $\dim \Sigma_A W < (n+1)q-1$ then $\sigma \circ f \simeq \omega_n$ for any homotopy section σ of g_n .*

Proof. Recall that g_i is the $(i+1)$ -fold join of ι_X . Thus by Mather (1976), Theorem 47, we obtain that, for each $i \geq 0$, $g_i: G_i \rightarrow X$ is $(i+1)q-1$ -connected. As g_i and

References

η_i^W have the same homotopy fibre, the Five lemma implies that $\eta_i^W: H_i \rightarrow \Sigma_A W$ is $(i+1)q-1$ -connected, too. By Whitehead (1978), Theorem IV.7.16, this means that for every CW-complex K with $\dim K < (i+1)q-1$, η_i^W induces a one-to-one correspondence $[K, H_i] \rightarrow [K, \Sigma_A W]$. Apply this to $K = \Sigma_A W$ and $i = n$: Since θ_n^W and $\bar{\sigma}$ are both homotopy sections of η_n^W , we obtain $\theta_n^W \simeq \bar{\sigma}$, and Proposition 6 implies the desired result. \square

Example 9 – Let $A = *$ and $W = S^{r-1}$, so $\Sigma_A W = S^r$, and $X = S^m$. In this case $\text{secat } \iota_X = \text{cat } S^m = 1$. Hence Theorem 2 means that if $r < 2m-1$, we have $\sigma \circ f \simeq \omega_1$, whatever can be f and $\sigma: X \rightarrow G_1(\iota_X)$, so $\text{hcat } f = 1$ and we get by Proposition 3 that the homotopy cofibre C of f has $\text{cat } C \leq 1$. (Notice that if $r < m$ then f is a nullhomotopic, so C is simply $S^m \vee S^{r+1}$.)

Example 10 – Let $A = *$, $\Sigma W \simeq \Sigma(S^{r-1} \vee S^{r-1}) \simeq S^r \vee S^r$, $X \simeq S^r \times S^r$ and consider $t_1: S^r \vee S^r \rightarrow S^r \times S^r$. Here $\text{secat } (\iota_X) = \text{cat } (S^r \times S^r) = 2$. For any $r \geq 1$, we have $\dim(S^r \vee S^r) = r < (2+1)r-1$, so $\text{hcat } (t_1) = 2$.

References

- Doeraene, J.-P. (2016). “Sectional Category of the Ganea Fibrations and Higher Relative Category”. *Chinese Journal of Mathematics*. doi: 10.1155/2016/8320742 (cit. on p. 190).
- Doeraene, J.-P. and M. El Haouari (2013). “Up-to-one approximations for sectional category and topological complexity”. *Topology and its Appl.* **160**, pp. 766–783 (cit. on pp. 189, 190, 194, 195, 197, 198).
- Iwase, N. (1998). “Ganea’s conjecture on Lusternik-Schnirelmann category”. *Bull. Lond. Math. Soc.* **30**, pp. 623–634 (cit. on p. 187).
- Mather, M. (1976). “Pull-backs in homotopy theory”. *Canad. Journ. Math.* **28** (2), pp. 225–263. ISSN: 0008-414X (cit. on pp. 188, 198).
- Whitehead, G. W. (1978). *Elements of homotopy theory*. **64**. Graduate texts in mathematics. New York: Springer-Verlag, pp. xxi+744 (cit. on p. 199).

Contents

1	The Hopf category	188
2	Hopf invariant and homotopy pushout	191
3	The Strong Hopf category	194
4	Equivalent conditions to get the Hopf category	198
	References	199
	Contents	i